

# Tips for a Good Colonoscopy Preparation-Plenvu

## Morning appointment

-Make sure you receive your colonoscopy prep and the appropriate instructions well before your procedure date, and read them completely as soon as you get them. It may be helpful to have a bottle of Miralax on hand in case needed to assist your bowel prep medication and prevent additional trips to the store once starting the prep process.

-Arrange for the time and privacy you need to complete the prep with as little stress as possible. Clear your schedule, and be at home on time to start your prep. If you have children or aging parents who need your attention, have someone else be available to them while you're indisposed.

-Water can get boring, so keep a variety of clear liquids on hand. On the day before your colonoscopy (when you're restricted to clear liquids) you can have popsicles, Jell-o, clear broth, coffee or tea (without milk or creamer), soft drinks, Italian ice or Gatorade. But take nothing with red, blue or purple dye. Drink extra liquids before and during your bowel prep, as well as after your colonoscopy. Hydration is important but you'll need to stop all liquids (including water) 4 hours prior to your exam time.

-Wear loose clothing and stay near the bathroom. Better yet, once the preparation starts to work, stay **IN** the bathroom (because when the urge hits, it's hard to hold back). Since you will be completely cleansing your bowel consider using baby wipes or a moisture barrier cream (similar to Aquafor) for comfort.

-The clear liquid diet and preparation are not a pleasant way to spend a day but your exam is only as good as your preparation. You should expect your output to move from solid to dark liquid to clear or clear yellow liquid by the end of your preparation process. Please make a note of the progress as you will be asked about your prep quality in pre-op. A poor preparation may result in missed lesions and could result in you having to repeat your procedure much sooner than would otherwise be needed. A good preparation will maximize our ability to locate and remove any pre-malignant lesion and will ensure that we give you the longest possible interval before needing to repeat your exam!


-Pay close attention to the diet and medication instructions noted below. You may need to make medication or diet changes up to 7 DAYS prior to your procedure. Medications such as: GLP-1, insulin, oral diabetic medications and blood thinning medications (other than aspirin) will likely have adjusted dosing schedules which will affect whether or not your procedure can be performed. Please make sure you have addressed this with your GI doctor and/or your prescribing doctor at least 1 week prior to your procedure; if not, your procedure will be cancelled.

See below:

**Daily GLP-1 medications should be held the day prior to your procedure. Weekly GLP-1 dosing should be held 7days prior to your procedure. SGLT2 drugs like; Jardiance (Empagliflozin), Farxiga (Dapagliflozin), Invokana (Canagliflozin) and Steglatro (Ertugliflozin) should be held 3 days prior to your procedure.**

# COLONOSCOPY TIMELINE

## Morning appointment



# PLENVU PREPARATION INSTRUCTIONS FOR COLONOSCOPY-Morning

## Appointment

### GENERAL INSTRUCTIONS:

- Expect to receive text invitations from our portal 2 weeks prior to your appointment (or at time of scheduling). We use this to communicate key instructions and to update your health history and assist anesthesia in preparation for your visit. Please complete the pre-procedure questionnaire located in the portal early to prevent receiving multiple phone calls from us.
- Expect a text or a phone call from our business office in the days prior to your appointment to review insurance information.
- Follow THESE instructions carefully, NOT the ones inside the prep box.
- Follow instructions given to you about any changes to your daily medications.
- **Pay close attention to the diet and medication instructions noted below.** You may need to make medication or diet changes up to 7 DAYS prior to your procedure. Medications such as: GLP-1, insulin, oral diabetic medications and blood thinning medications (other than aspirin) will likely have adjusted dosing schedules which will affect whether or not your procedure can be performed. Please make sure you have addressed this with your GI doctor and/or your prescribing doctor at least 1 week prior to your procedure; if not, your procedure may be canceled.
- You should NOT have ANYTHING FOUR HOURS prior to your exam. Including gum, mints and ice chips.
- **If you do not follow these directions your colonoscopy may be canceled.**

INSURANCE NOTIFICATION:- If precertification is required, it will be obtained by our Central Billing Office. Contact your insurance company prior to the procedure to make sure this has been done.

- If you have changed insurance since the last time you have been seen, you must notify us of this change.
- There will be a facility and anesthesia charge in addition to the physician's fee.
- **Please bring your insurance card and driver's license to the procedure.**

### 5-7 DAYS BEFORE THE EXAM:

- You will likely have to stop BLOOD THINNERS prior to your exam. Your doctor will tell you how long you will need to be off the blood thinners before and after the exam. Do NOT stop your aspirin unless otherwise instructed.
- Discontinue fiber supplements and any medications containing iron (Metamucil, Citrucel, multivitamins with iron, etc.).
- IF YOU ARE DIABETIC, contact your primary care physician for instructions on how to take your medications on the day of the preparation and the procedure day.
- **If you take any type of GLP-1 agonist medication such as Ozempic/Wegovy (Semaglutide), Trulicity (Dulaglutide), Zepbound or Mounjaro (Trizeptide) or SGLT2 medications such as Jardiance (Empagliflozin), Farxiga (Dapagliflozin), Invokana (Canagliflozin) and Steglatro (Ertugliflozin) be sure to follow the hold instructions above. For you safety, if hold instructions not followed, your procedure will be cancelled.**

### 3 DAYS BEFORE THE EXAM:

- Stop eating all high fiber foods: No raw fruits or vegetable, whole grain wheat or multi-grain foods, nuts, popcorn, seeds, bran, bulking agents.

For insurance questions please contact the Lone Star Insurance team at 817-402-7700      See back→

## THE DAY BEFORE THE EXAM:

-Drink only **CLEAR LIQUIDS** for breakfast, lunch, and dinner. Start drinking liquids in the morning, and drink all day. Do not just drink the liquids recommended around the time of the actual preparation. The more you drink all day, the better the prep will be.

- NO solids, or milk products.

- NO alcohol on the day before or the day of the procedure.

- CLEAR LIQUIDS are: Strained fruit juices without pulp (apple, lemonade, white grape) Water, Gatorade, popsicles, snow cones, carbonated soft drinks, Jell-O, Clear broth and bouillon, Coffee or tea WITHOUT creamer/milk. AVOID RED, BLUE and PURPLE LIQUIDS

*Ondansetron (Zofran) prescription-If your bowel prep kit also came with an RX for Zofran please take 1-2 pills by mouth 30mins-1hrs before or after each bowel prep dose as needed to prevent nausea.*

### **Please start the first dose of prep at 6:00pm**

1. Using the mixing container, mix the contents of the Dose 1 pouch (Mango flavor) with at least 16 ounces of water by shaking or using a spoon until it's completely dissolved. This may take up to 2-3 minutes. Take your time – slowly finish the dose within 30 minutes.
2. Refill the container with at least 16 ounces of clear liquid. Again take you time and slowly finish all of it within 30 minutes.

**\*\*If no BM 2 hours after the first dose (of ANY bowel prep), then take 1-2 doses of Miralax in 8oz of water**

### **Please take your second dose of prep 5 hours prior to arrival time**

1. Using the mixing container, mix the contents of Dose 2 (Pouch A and Pouch B, Fruit Punch flavor) with at least 16 ounces of water by shaking or using a spoon until it's completely dissolved. This may take up to 2-3 minutes. Take your time – slowly finish the dose within 30 minutes.
2. Refill the container with at least 16 ounces of clear liquid. Again, take your time and slowly finish all of it within 30 minutes.
3. **Nothing more by mouth within 4 hours of your exam time. This includes chewing gum, hard candy, water and chewing tobacco. Doing any of these will delay and possibly cancel your procedure.**

## DAY OF EXAM:

-If you are on heart or blood pressure medications, take them with a sip of water 2 hours or more before the exam. Other medications may be taken as directed by doctor or pre-op nurse.

-Otherwise, **NOTHING BY MOUTH**, within 4 hours of the exam. The second dose of prep should be finished 4 hours **prior** to arrival.

-You will be in recovery for 30-45 minutes after the exam. Average total time at the facility is around 2 hours.

-You will need a friend or family member to drive you home. You should **NOT** drive, work, exercise, or drink alcohol for the rest of the day after your exam.

PLEASE CALL TDDC OFFICE DIRECTLY IF YOU ARE UNABLE TO COMPLETE YOUR PREP OR FEEL THAT YOU ARE NOT CLEAR AFTER COMPLETING THE PREP.

**Texas Digestive Disease Consultants-24 HOURS CALL**

**Southlake (817)424-1525**

**Alliance (817)741-4347**

**Flower Mound (972)691-3777**